

Land use Changes and Northern Hemisphere Cooling

B. Govindasamy, P. B. Duffy and K. Caldeira

Climate and Carbon Cycle Modeling group Atmospheric Science
Division Lawrence Livermore National Laboratory
Livermore, CA

Abstract. Recent reconstructions of Northern Hemisphere mean temperatures over the past millennium show a long-term cooling of about 0.25K between 1000 and 1900 AD, prior to the 20th century's warming. In this paper, we present the results of equilibrium climate model simulations that indicate that the land-use change occurring over this period may largely explain this observed cooling, although other factors also could have played a significant role. The simulated annual mean cooling due to land-use change is 0.25K globally and 0.37 K for the Northern Hemisphere, suggesting that the cooling of the prior centuries could have been largely the result of anthropogenic interference in the climate system.

Introduction

Recent reconstructions of Northern Hemisphere temperatures over the past millennium show a long-term cooling of about 0.25 K between 1000 and 1900 AD, prior to 20th century's warming (Mann *et al.*, 1998; 1999). It has been suggested that this observed cooling is related to changes in astronomical forcing (Berger, 1988). An alternate hypothesis is that significant century-scale variability may be associated with solar irradiance variations (Lean *et al.*, 1995). Here, we present the results of climate model simulations that indicate that the cooling due to land-use change occurring over this period is of the sign and magnitude to explain most of this observed cooling.

Previous studies of climate change and land use change have mostly looked at the effects of tropical deforestations and particularly the Amazonian deforestation (Shukla *et al.*, 1990; Nobre *et al.*, 1991; Dickinson and Kennedy, 1992; Sellers *et al.*, 1993). These studies show that changes in land use such as conversion of forest to grassland increase surface albedo and decrease surface roughness. The change in albedo decreases the net solar radiation absorbed and results in decreased surface temperature. The change in surface roughness decreases evaporation and leads to increased surface temperature. Both changes result in decreased moisture convergence and reduced convective precipitation due to increased static stability (albedo effect) or due to reduced evaporation (roughness effect). These studies reveal that the change in surface roughness increases the surface temperatures more than they were decreased by the change in albedo. Costa and Foley (2000) find that the effects of deforestation and increasing CO₂ on precipitation tend to counteract one another but both tend to warm the Amazon basin.

Boreal forest ecosystems may also affect climate (Bonan *et al.*, 1992; Foley *et al.*, 1994; Bonan, 1997; Hansen *et al.*, 1999; Chase *et al.*, 2000). The increase in albedo caused by deforestation of boreal forest apparently overwhelms the warming effect due to reduced evaporation. Further, in mid and high latitudes, deforestation also exposes any underlying snow, reinforcing the cooling in a 'snow-albedo' feedback. Vegetation and soil feedbacks have been implicated in climate change (Foley *et al.*, 1994; Kutzbach *et al.*, 1996; Coe and Bonan, 1997) in the early Holocene.

Since most of the land use change took place in mid and high latitudes before the industrial era, it is possible that conversion of natural forests into cropland causes cooling and that this can explain much of the observed Northern Hemisphere cooling in the last millennium. In this paper, we use an atmospheric general circulation model (AGCM) coupled to a simplified model of the ocean and sea ice, to investigate the climate change due to the replacement of natural vegetation into croplands that occurred prior to the industrial era. Indeed, we find that such a land use change causes a decrease in simulated Northern Hemisphere and global mean temperatures similar to that estimated from paleo-data reconstructions. Chase *et al.* (2000) investigated the same issue of impacts of land-use change using the same AGCM and datasets, but their study was limited by the use of prescribed sea surface temperatures.

The Model

We adopted Version 3 of the Community Climate Model (CCM3) developed at the National Center for Atmospheric Research (Kiehl *et al.*, 1998). The horizontal resolution is approximately 2.8° in latitude and 2.8° in longitude. The model has 18 levels in the vertical. An important aspect of CCM3 is that it has very little systematic bias in the top-of-atmosphere and surface energy budgets. Its climatology is in reasonable agreement with observations (Kiehl *et al.*, 1998). We adopted a version of the model with a simple slab ocean-thermodynamic sea ice model, which allows for a simple interactive surface for the ocean and sea ice components of the climate system. The slab ocean model employs a spatially and seasonally prescribed ocean heat transport and spatially prescribed mixed layer-depth, which ensures replication of realistic sea surface temperatures and ice distributions for the present climate.

Experiments

We performed two equilibrium climate simulations. One of these simulations, "1000AD", used potential natural vegetation (Chase *et al.*, 1996; 2000); the other simulation, "1900AD", used the standard CCM current vegetation dataset (Bonan, 1996). For CO₂ and other trace gases such as CH₄,

Copyright 2001 by the American Geophysical Union.

Paper number 2000GL006121.
0094-8276/01/2000GL006121\$05.00




Figure 1. Change in annual mean exposed Leaf Area Index (top panel) and net solar radiation at the top of the atmosphere (bottom panel) due to the conversion of natural potential vegetation into croplands by anthropogenic interference prior to the industrial era.

N_2O , CFC-11 and CFC-12, the concentrations were specified at pre-industrial levels in both simulations. For both simulations, the model was run for 25 years from an initial state corresponding to pre-industrial climate. The climate statistics presented are the mean values over the last 15 years of the model simulations.

The natural potential vegetation cover is created by taking the standard CCM dataset and filling in agricultural areas with appropriate natural vegetation types in keeping with standard CCM3 vegetation classifications (Chase *et al.*, 2000). The natural potential vegetation type is estimated from the relationship between potential Leaf Area Index (LAI) and vegetation types (Neilson and Marks, 1994). Potential LAI is estimated using long-term precipitation, temperature, and soil data and an empirical relationship between transpiration and LAI (Nemani and Running, 1989). The use of standard CCM3 vegetation types that correspond most closely to the estimated vegetation types (Nemani and Running, 1989) may contribute some inaccuracy but is unlikely to affect our basic results.




Figure 2. a) Simulated surface temperature change due to conversion of natural vegetation to croplands. Simulated surface temperature cools most significantly over central and eastern North America, Middle East, and Southeast Asia. The contour interval is 0.5 K. b) The regions where the cooling is significant at the 5 % level.

Changes in exposed LAI and net solar radiation at the top of the atmosphere in regions of land use change are shown in Fig. 1. The affected regions are mostly in Europe, North America, India and Southeast Asia. Savanna, various types of forests, grassland and shrubland were replaced by croplands. Such a conversion generally leads to a reduction in mean net solar radiation at the surface and top of the atmosphere. Our radiative forcing estimations yielded a mean shortwave radiative forcing of -0.28 W m^{-2} over land area due to these changes.

Results

The simulated annual mean cooling due to land-use change is 0.25 K globally, 0.37 K for the Northern Hemisphere and 0.41 K over land. We notice a corresponding decrease in precipitable water and increase in sea ice volumes

Table 1. Climate Statistics: Annual means of global, Northern Hemisphere and Land means of surface temperature, precipitable water and sea ice volume in "1000AD" and "1900AD" simulations and the difference between them.

Case	Surface Temperature (K)			Precipitable water (mm)			Sea ice Vol. ($\times 10^{12} \text{ m}^3$)	
	Global	NH	Land	Global	NH	Land	Global	NH
1000AD	285.75	286.20	278.43	25.32	26.05	19.52	50.65	38.54
1900AD	285.50	285.83	278.02	24.87	25.16	19.21	51.20	39.43
Change	-0.25	-0.37	-0.41	-0.45	-0.89	-0.31	0.55	0.89

Table 2. Climate Statistics: Annual means of global, Northern and Southern Hemisphere and Land means net surface solar radiation, precipitation, and evaporation in the "1000AD" and "1900AD" simulations and the difference between them.

Case	Net surface solar radiation (W m ⁻²)			Precipitation (mm day ⁻¹)			Evaporation (mm day ⁻¹)		
	1000AD	1900AD	Change	1000AD	1900AD	Change	1000AD	1900AD	Change
Global	172.24	171.77	-0.47	2.98	2.98	0.0	2.98	2.98	0.0
NH	170.56	170.13	-0.43	3.03	2.93	-0.1	2.87	2.87	0.0
SH	173.92	173.41	-0.51	2.93	3.03	0.1	3.09	3.09	0.0
Land	145.77	143.91	-1.86	2.16	2.21	0.05	1.47	1.50	0.03

in the current vegetation case (Table 1). Our results are in qualitative agreement with a recent study (Hansen *et al.*, 1999) where a simulated global mean cooling of 0.14 K is obtained.

The model predicts that eastern and central North America, Middle East, and Southeast Asia cool more than the Northern Hemisphere average (Fig. 2a). The centers of cooling are located in these regions and extend to Greenland, Europe, Northern Africa and India. The cooling in these regions is statistically significant at the 5% level (Fig. 2b). In contrast, areas with warming and all temperature anomalies in the high latitudes are generally not statistically significant at this level.

In order to understand this simulated cooling, we show in Table 2 the annual means of global, Northern and Southern Hemisphere and land surface means of net surface solar radiation, precipitation and surface evaporation. We notice a decrease in net solar radiation in the global and Northern Hemisphere in the current vegetation case. This decrease is in qualitative agreement with a recent modeling study (Hansen *et al.*, 1999). Over land, a decrease of 1.86 W m⁻² in the net absorbed solar radiation at the surface is simulated. Therefore, we conclude that the simulated cooling occurs primarily as a result of the higher albedo of agroecosystems as compared with the natural ecosystems they replaced.

In tropical deforestation studies (Shukla *et al.*, 1990; Nobre *et al.*, 1991; Dickinson and Kennedy, 1992; Sellers *et al.*, 1993), it was found that the conversion of tropical forests into croplands significantly affected mean precipitation and evaporation due to local changes in tropical convection. In contrast to these studies, we find little changes in mean precipitation and evaporation (Table 2). This is because, in our study, the deforestation occurred mostly in mid latitudes where precipitation dynamics is less affected by local convection and is instead dominated by large-scale condensation associated with baroclinically driven storm tracks. Therefore, we conclude that little change in the global scale hydrologic cycle is associated with historical land-use changes.

Discussion

We have performed "equilibrium" climate simulations using an atmospheric general circulation model coupled to a slab-ocean and thermodynamic sea-ice model (Kiehl *et al.*, 1998). It is unlikely that the transition between "natural" and "modern" vegetation was smooth (Ramankutty and Foley, 1998; 1999). It is possible that other GCMs would yield quantitatively different results, because the results may be highly sensitive to the formulation of the model and the

parameterization of various physical processes. Simulations using a coupled atmosphere, dynamic sea ice and ocean general circulation model would include dynamical feedbacks that could amplify the regional or global climate change. Nonetheless, our results indicate that deforestation can cause significant climate perturbations and further efforts should be made to improve our understanding of the processes involved.

Our results indicate that land use change could explain most of the cooling inferred for the period from 1000 to 1900 AD (Mann *et al.*, 1998; 1999). However, other factors could have also contributed to this cooling, including internal oscillations related to ocean-atmosphere coupling, changes in earth's orbital parameters (Berger, 1988), or century-scale climate variability associated with solar irradiance variations (Lean *et al.*, 1995). We have verified that our model is relatively insensitive to changes in earth's orbital parameters. The change in global and annual mean surface temperature is of the order of 0.01 K for orbital parameters corresponding to glacial (114,000 BP) and interglacial (125,000 BP) periods. The maximum range of changes in solar irradiance is ~ 4 W m⁻² from 1600 AD to present (Lean and Rind, 1998). Taking into consideration Earth's albedo and geometry, this corresponds to a climate forcing of 0.7 W m⁻², and a temperature increase of 0.32 K, since the climate sensitivity of our model is 0.45 K/Wm⁻² (Govindasamy and Caldeira, 2000). This is indeed similar in magnitude but opposite in sign to the temperature change we find due to land use change. Because solar luminosity is inferred to have increased since 1600 AD, solar variability may be an important factor affecting climate over the past millenium, but is unlikely to explain the cooling between in the period prior to 1900 AD.

Nevertheless, we wish to emphasize that the goal of this paper not to explain all of the factors that could have contributed to the inferred Northern Hemisphere cooling since 1000 AD, or to systematically investigate all the possible causes of this cooling. Rather, our goal is to assess the climatic impact of land use changes and understand whether this could have been an important factor contributing to Northern Hemisphere cooling. Since the model simulated temperature change is similar in sign and magnitude to the observed temperature change since 1000 AD, we suggest that land use change is a potentially important factor contributing to this cooling. Of course, other factors may also prove to be important. The observed datasets are too sparse at present to test the land-use-change/global-cooling hypothesis with detailed spatial resolution; however, this prediction is in principle testable. We urge the collection of additional paleoclimate datasets that can be used to test this hypothesis that land-use changes contributed to the observed Northern Hemisphere cooling between 1000 and 1900 AD.

Acknowledgements. This work was performed under the auspices of the U.S. Department of Energy by the Lawrence Livermore National Laboratory under contract No. W-7405-Eng-48. We are thankful to Dr. Thomas Chase of Colorado State University for providing the potential natural vegetation data.

References

- Berger, A., Milankovitch theory and climate, *Rev. of Geophys.*, **26**, 624-657, 1988.
- Bonan, G. B., D. Pollard, and S. L. Thompson, Effects of boreal forest vegetation on global climate, *Nature*, **359**, 716-718, 1992.
- Bonan, G. B., A land surface model for ecological, hydrological, and atmospheric studies: Technical description and user's guide, NCAR Tech. Note, *NCARTN-417+STR*, 1996.
- Bonan, G. B., Effects of land use on the climate of the United States, *Climate Change*, **37**, 449-486, 1997.
- Chase, T., R. A. Pielke, T. G. F. Kittel, R. Nemani & S. W. Running, Sensitivity of a general circulation model to global changes in leaf area index, *J. Geophys. Res.*, **101**, 7393-7408, 1996.
- Chase, T. N., R. A. Pielke, T. G. F. Kittel, R. R. Nemani and S. W. Running, Simulated impacts of historical land cover changes on global climate in northern winter, *Climate Dynamics*, **16**, 93-105, 2000.
- Coe, M. T., and G. B. Bonan, Feedbacks between climate and surface water in northern Africa during the middle Holocene, *J. Geophys. Res.*, **102**, 11087-11101, 1997.
- Costa, M. H., and J. Foley, Combined effects of deforestation and doubled atmospheric CO₂ concentration on the climate of Amazonia, *J. Climate*, **13**, 18-34, 2000.
- Dickinson, R. E., and P. Kennedy, Impacts on regional climate of Amazon deforestation, *Geophys. Res. Lett.*, **19**, 1947-1950, 1992.
- Foley, J. A., J. E. Kutzbach, M. T. Coe, and S. Levis, Feedbacks between climate and boreal forests during the Holocene epoch, *Nature*, **371**, 52-54, 1994.
- Govindasamy, B., and K. Caldeira, Geoengineering Earth's radiation balance to mitigate CO₂-induced climate change, *Geophys. Res. Lett.*, **27**, 2141-2144, 2000.
- Hansen, J. E., M. Sato, A. Lacis, R. Ruedy, I. Tegen & E. Mathews, Climate forcings in the industrial era, *Proc. Natl. Acad. Sci.*, **95**, 12753-12758, 1999.
- Kiehl, J. T., J.J.Hack, G.B.Bonan, B.Y.Boville, B.P.Briegleb, D.L.Williamson, and P.J.Rasch, The National Center for Atmospheric Research Community Climate Model: CCM3, *J. Climate*, **11**, 1131-1149, 1998.
- Kutzbach, J., G. Bonan, J. Foley, and S. P. Harrison, Vegetation and soil feedbacks on the response of the African monsoon to orbital forcing in the early to middle Holocene, *Nature*, **384**, 623-626, 1996.
- Lean, J., J. Beer & R. Bradley, Reconstruction of solar irradiance since 1610: Implications for climate change, *Geophys. Res. Lett.*, **22**, 3195-3198, 1995.
- Lean, J., and D. Rind, Climate forcing by changing solar radiation, *J. Climate*, **11**, 3069-3094, 1998.
- Mann, M. E., R. S. Bradley & M. K. Hughes, Global-scale temperature patterns and climate forcing over the past six centuries, *Nature*, **392**, 779-787, 1998.
- Mann, M. E., R. S. Bradley & M. K. Hughes, Northern Hemisphere temperatures during the past millennium: Inferences, uncertainties and limitations, *Geophys. Res. Lett.*, **26**, 759-762, 1999.
- Neilson, R. P., and D. Marks, A global perspective of regional vegetation and hydrologic sensitivities from climate change, *J. Vegetation Sci.*, **5**, 715-730, 1994.
- Nemani, R.R., and S. W. Running, Testing a theoretical climate-soil-leaf area hydrologic equilibrium of forests using a satellite data and ecosystem simulation, *Agric. Forest Meteorol.*, **44**, 245-260, 1989.
- Nobre, C. A., P. J. Sellers, J. Shukla, Amazonian deforestation and regional climate change, *J. climate*, **4**, 957-988, 1991.
- Ramankutty, N., and J. A. Foley, Characterizing patterns of global land use: An analysis of global croplands data, *Global Biogeochemical Cycles*, **12**, 667-685, 1998.
- Ramankutty, N., and J. A. Foley, Estimating historical changes in global land cover: Croplands from 1700 to 1992, *Global Biogeochemical Cycles*, **13**, 997-1027, 1999.
- Sellers, H. A., R. E. Dickinson, T. B. Durbidge, P. J. Kennedy, K. McGuffie, and A. J. Pitman, Tropical deforestation - Modeling local-scale to regional-scale climate change, *J. Geophys. Res.*, **98**, 7289-7315, 1993.
- Shukla, J., C. Nobre, and P. J. Sellers, Amazon deforestation and climate change, *Science*, **247**, 1322-1325, 1990.
- B. Govindasamy, P. B. Duffy and K. Caldeira, Climate and Carbon Cycle Group, Lawrence Livermore National Laboratory, 7000 East Ave., L-103, Livermore, CA 94550. (bala@llnl.gov; pduffy@llnl.gov; kenc@llnl.gov)

(Received May 3, 2000; revised August 8, 2000; accepted October 5, 2000.)